

Species that have bred at NC Zoo

Amethyst Starling

Bali Mynah

Blue-crowned Laughing Thrush

Blue-grey Tanager

Common (white-rumped) Shama Thrush

Fairy Bluebird

Golden Crested Mynah

Golden-headed Manakin

Golden White-eye

Hooded Pitta

Lesser Green Broadbill

Pekin Robin (Red-billed Leiothrix)

Red-capped Cardinal

Red-faced Liocichla

Snowy-headed Robin Chat

Turquoise Tanager

Yellow-rumped Cacique

- African Pygmy Goose
- Blue-crowned Hanging Parrot
- Crested Wood Partridge
- •Green Woodhoopoe
- Hottentot Teal
- Nicobar Pigeon
- Sunbittern
- Victoria-crowned Pigeon
- Wattled Cranes
- White-winged Wood Duck



NC ZOO AVIARY


- Opened in 1982
- 400,000 cu ft²
- 138' Diameter
- 55' High
- ~35 Species
- ~110 Birds
- 2000 Plants







Natural Habitat



- Ficus trees
- Macadamia nut tree
- Caricature plant
- Banyan tree
- Flame vine
- Guyana chestnut tree
- Stilt root palm




















Birds use materials found in exhibit to build nests



Problems we have had in the past with letting birds nest in our large free flight aviary

- Dietary concerns
- Parasites
- Allowing chicks to fledge in exhibit
- Moving nest too close to fledge date



Diet

Unlimited waxworms

Insect diet

Supplements

Ingredient change



Parasites

Parents parasite load

Nematodes

Prophylactic treatment

Fledging in exhibit

- Dangers:
 - Pool
 - -Visitors
 - –Small crevices/stuck in plants
 - -Other species interfering
 - Parents uncomfortable going to ground

Led to high mortality in fledglings



Moving nest too close to fledge date

Fledging early

 Two stressful events in a short period of time (Move and fledge)

Led to high chick mortality

 Monitoring nesting events by close observation of nest construction, incubation, and suspected hatches within the exhibit.





Relocate Family to Trap Cage 5 Days After Hatch

- Trap Adults First
- Cut Down Nest With Chicks and Relocate to Trap Cage with Adults



- If Nest Not
 Accessible, Chicks
 Allowed to Fledge Out
 on Exhibit
 - Place Fledglings in Small Mesh Cage
 - Trapping Parents





- Pekin Robin Family Diet
 - Marion Red Apple
 Paradise Pellet
 - Mazuri Bird Gel
 - Mazuri Small Bird
 Particle Breeder
 - Fruit and Veg Mix




- Pekin Robin Family Diet: Protein
 - Headless Male Crickets
 - Pinkie Pieces
 - Unlimited mealworms
 - Limited waxworms and superworms

- Family Moved to
 Avian Propagation

 11 Days After Chicks
 Fledge
- At Prop, Juveniles
 Weaned and
 Separated From
 Adults



Red-faced Liocichla





















Golden White-eye




Yellow-rumped Cacique



Golden-crested Mynah




Avian Propagation



Avian Propagation







Avian Propagation



- Nesting Materials
 Offered
 - Raffia
 - Palm Stalks with Fibers
 - Grassy Goodnight

